

Section 2.2: Solving Systems of Equations by Substitution

Objective: Solve systems of equations using substitution.

Solving a system by graphing has several limitations. First, it requires the graph to be perfectly drawn. If the lines are not straight we may arrive at the wrong answer. Second, graphing is not a great method to use if the answer is really large, such as $(100, -75)$, or if the answer contains a decimal that the graph will not help us find, such as $(3.2134, 2.17)$. For these reasons we will rarely use graphing to solve our systems. Instead, an algebraic approach will be used.

The first algebraic approach is called substitution. We will build the concepts of substitution through several examples, then end with a five-step process to solve problems using this method.

Example 1.

Solve the systems of equations by using substitution:

$$\begin{aligned}x &= 5 \\ y &= 2x - 3\end{aligned}$$

$x = 5$	We already know $x = 5$, substitute this into
$y = 2x - 3$	the other equation
$y = 2(\mathbf{5}) - 3$	Evaluate, multiply first
$y = 10 - 3$	Subtract
$y = 7$	We now also have y
$(5, 7)$	Our Solution

When we know what one variable equals we can plug that value (or expression) in for the variable in the other equation. It is very important that when we substitute, the substituted value goes in parentheses. The reason for this is shown in the next example.

Example 2.

Solve the systems of equations by using substitution:

$$\begin{aligned}2x - 3y &= 7 \\ y &= 3x - 7\end{aligned}$$

$2x - 3y = 7$	We know $y = 3x - 7$; substitute this into the
$y = 3x - 7$	other equation
$2x - 3(\mathbf{3x - 7}) = 7$	Solve this equation, distributing -3 first

$$\begin{array}{r}
2x - 9x + 21 = 7 \quad \text{Combine like terms } 2x - 9x \\
-7x + 21 = 7 \quad \text{Subtract 21} \\
-7x + 21 = 7 \\
\hline
-21 \quad -21 \\
-7x = -14 \\
\hline
\frac{-7x}{-7} = \frac{-14}{-7} \quad \text{Divide by } -7 \\
x = 2 \quad \text{We now have our } x; \text{ plug into the } y = \\
\text{equation to find } y \\
y = 3(2) - 7 \quad \text{Evaluate, multiply first} \\
y = 6 - 7 \quad \text{Subtract} \\
y = -1 \quad \text{We now also have } y \\
(2, -1) \quad \text{Our Solution}
\end{array}$$

By using the entire expression $3x - 7$ to replace y in the other equation we were able to reduce the system to a single linear equation, which we can easily solve for our first variable. However, the lone variable (a variable without a coefficient) is not always alone on one side of the equation. If this happens we can isolate it by solving for the lone variable.

Example 3.

Solve the systems of equations by using substitution: $3x + 2y = 1$
 $x - 5y = 6$

$$\begin{array}{r}
3x + 2y = 1 \quad \text{Lone variable is } x; \text{ isolate by adding } 5y \text{ to both sides.} \\
x - 5y = 6 \\
\hline
+5y \quad +5y \\
x = 6 + 5y \quad \text{Substitute this into the untouched equation} \\
3(6 + 5y) + 2y = 1 \quad \text{Solve this equation, distributing 3 first} \\
18 + 15y + 2y = 1 \quad \text{Combine like terms } 15y + 2y \\
18 + 17y = 1 \\
\hline
-18 \quad -18 \\
17y = -17 \\
\hline
\frac{17y}{17} = \frac{-17}{17} \quad \text{Divide both sides by 17} \\
y = -1 \quad \text{We have our } y; \text{ plug this into the } x = \text{ equation to find } x \\
x = 6 + 5(-1) \quad \text{Evaluate, multiply first} \\
x = 6 - 5 \quad \text{Subtract} \\
x = 1 \quad \text{We now also have } x \\
(1, -1) \quad \text{Our Solution}
\end{array}$$

The process in the previous example is how we will solve problems using substitution. This process is described and illustrated in the following table which lists the five steps to solving by substitution.

Problem	$4x - 2y = 2$ $2x + y = -5$
1. Find the lone variable	Second Equation, y $2x + y = -5$
2. Solve for the lone variable	$-2x \quad -2x$ $y = -5 - 2x$
3. Substitute into the untouched equation	$4x - 2(-5 - 2x) = 2$
4. Solve	$4x + 10 + 4x = 2$ $8x + 10 = 2$ $\quad -10 \quad -10$ $\frac{8x}{8} = \frac{-8}{8}$ $x = -1$
5. Plug into lone variable equation and evaluate	$y = -5 - 2(-1)$ $y = -5 + 2$ $y = -3$
Solution	$(-1, -3)$

Sometimes we have several lone variables in a problem. In this case we will have the choice on which lone variable we wish to solve for; either choice will give the same final result.

Example 4.

Solve the system of equations by using substitution.

$$\begin{array}{ll}
 x + y = 5 & \text{Find the lone variable: } x \text{ or } y \text{ in first, or } x \text{ in second.} \\
 x - y = -1 & \text{We will chose } x \text{ in the first} \\
 x + y = 5 & \text{Solve for the lone variable; subtract } y \text{ from both sides} \\
 \underline{-y \quad -y} & \\
 x = 5 - y & \text{Plug into the untouched equation, the second equation} \\
 (5 - y) - y = -1 & \text{Solve (parentheses are not needed here); combine like} \\
 & \text{terms} \\
 5 - 2y = -1 & \text{Subtract 5 from both sides} \\
 \underline{-5 \quad -5} & \\
 -2y = -6 & \\
 \underline{-2y = -6} & \\
 \underline{-2 \quad -2} & \text{Divide both sides by } -2
 \end{array}$$

$$\begin{array}{ll}
 y = 3 & \text{We have our } y! \\
 x = 5 - (3) & \text{Plug into lone variable equation; evaluate} \\
 x = 2 & \text{Now we have our } x \\
 (2, 3) & \text{Our Solution}
 \end{array}$$

Just as with graphing it is possible to have no solution \emptyset (parallel lines) or infinite solutions (same line) with the substitution method. While we won't have a parallel line or the same line to look at and conclude if it is one or the other, the process takes an interesting turn as shown in the following example.

Example 5.

$$\begin{array}{ll}
 y + 4 = 3x & \text{Find the lone variable, } y, \text{ in the first equation} \\
 2y - 6x = -8 & \\
 y + 4 = 3x & \text{Solve for the lone variable; subtract 4 from both sides} \\
 \quad -4 \quad -4 & \\
 \hline
 y = 3x - 4 & \text{Plug into untouched equation} \\
 2(\mathbf{3x - 4}) - 6x = -8 & \text{Solve; first distribute 2 through parentheses grouping} \\
 6x - 8 - 6x = -8 & \text{Combine like terms } 6x - 6x \\
 \quad -8 = -8 & \text{Variables are gone! A true statement.} \\
 \text{Infinite solutions} & \text{Our Solution}
 \end{array}$$

Because we had a true statement, and no variables, we know that anything that works in the first equation, will also work in the second equation. However, we do not always end up with a true statement.

Example 6.

$$\begin{array}{ll}
 6x - 3y = -9 & \text{Find the lone variable, } y, \text{ in the second equation} \\
 -2x + y = 5 & \\
 -2x + y = 5 & \text{Solve for the lone variable; add } 2x \text{ to both sides} \\
 \quad +2x \quad +2x & \\
 \hline
 y = 5 + 2x & \text{Plug into untouched equation} \\
 6x - 3(\mathbf{5 + 2x}) = -9 & \text{Solve, first distribute through parentheses grouping} \\
 6x - 15 - 6x = -9 & \text{Combine like terms } 6x - 6x \\
 \quad -15 = -9 & \text{Variables are gone! A false statement.} \\
 \text{No solution } \emptyset & \text{Our Solution}
 \end{array}$$

Because we had a false statement and no variables, we know that no numerical values will work in both equations.

World View Note: French mathematician Rene Descartes wrote a book which included an appendix on geometry. It was in this book that he suggested using letters from the end of the alphabet for unknown values. This is why often we are solving for the variables x , y and z .

One more question needs to be considered, what if there is no lone variable? If there is no lone variable substitution can still work to solve, we will just have to select one variable to solve for and use fractions as we solve.

Example 7.

$$\begin{array}{ll}
 5x - 6y = -14 & \text{No lone variable} \\
 -2x + 4y = 12 & \text{We will solve for } x \text{ in the first equation} \\
 5x - 6y = -14 & \text{Solve for our variable, add } 6y \text{ to both sides} \\
 \quad \quad \quad + 6y & + 6y \\
 \hline
 5x = \frac{-14}{5} + \frac{6y}{5} & \text{Divide each term by } 5 \\
 x = \frac{-14}{5} + \frac{6y}{5} & \text{Plug into untouched equation} \\
 -2\left(\frac{-14}{5} + \frac{6y}{5}\right) + 4y = 12 & \text{Solve, distribute through parenthesis} \\
 \quad \quad \quad \frac{28}{5} - \frac{12y}{5} + 4y = 12 & \text{Clear fractions by multiplying by } 5 \\
 \frac{28(5)}{5} - \frac{12y(5)}{5} + 4y(5) = 12(5) & \text{Reduce fractions and multiply} \\
 28 - 12y + 20y = 60 & \text{Combine like terms } -12y + 20y \\
 28 + 8y = 60 & \text{Subtract } 28 \text{ from both sides} \\
 \quad \quad \quad -28 & -28 \\
 \hline
 \quad \quad \quad 8y = 32 & \text{Divide both sides by } 8 \\
 \quad \quad \quad \frac{8y}{8} = \frac{32}{8} & \\
 \quad \quad \quad y = 4 & \text{We have our } y \\
 x = \frac{-14}{5} + \frac{6(4)}{5} & \text{Plug into lone variable equation, multiply} \\
 x = \frac{-14}{5} + \frac{24}{5} & \text{Add fractions} \\
 x = \frac{10}{5} & \text{Reduce fraction} \\
 x = 2 & \text{Now we have our } x \\
 (2, 4) & \text{Our Solution}
 \end{array}$$

Using the fractions could make the problem a bit trickier. This is why we have another method for solving systems of equations that will be discussed in another lesson.

2.2 Practice

Solve each system by substitution.

1) $y = -3x$
 $y = 6x - 9$

2) $y = x + 5$
 $y = -2x - 4$

3) $y = -2x - 9$
 $y = 2x - 1$

4) $y = -6x + 3$
 $y = 6x + 3$

5) $y = 6x + 4$
 $y = -3x - 5$

6) $y = 3x + 13$
 $y = -2x - 22$

7) $y = 3x + 2$
 $y = -3x + 8$

8) $y = -2x - 9$
 $y = -5x - 21$

9) $y = 2x - 3$
 $y = -2x + 9$

10) $y = 7x - 24$
 $y = -3x + 16$

11) $y = 6x - 6$
 $-3x - 3y = -24$

12) $-x + 3y = 12$
 $y = 6x + 21$

13) $y = -6$
 $3x - 6y = 30$

14) $6x - 4y = -8$
 $y = -6x + 2$

- 15) $y = -5$
 $3x + 4y = -17$
- 16) $7x + 2y = -7$
 $y = 5x + 5$
- 17) $-2x + 2y = 18$
 $y = 7x + 15$
- 18) $y = x + 4$
 $3x - 4y = -19$
- 19) $y = -8x + 19$
 $-x + 6y = 16$
- 20) $y = -2x + 8$
 $-7x - 6y = -8$
- 21) $7x - 3y = -7$
 $y = 4x + 1$
- 22) $3x - 2y = -13$
 $4x + 2y = 18$
- 23) $x - 5y = 7$
 $2x + 7y = -20$
- 24) $3x - 4y = 15$
 $7x + y = 4$
- 25) $-2x - 5y = -5$
 $x - 8y = -23$
- 26) $6x + 4y = 16$
 $-2x + 9y = -3$
- 27) $-6x + y = 20$
 $-3x - 3y = -18$
- 28) $7x + 5y = -13$
 $x - 4y = -16$
- 29) $3x + y = 9$
 $2x + 8y = -16$
- 30) $-5x - 5y = -20$
 $-2x + y = 7$

$$31) \begin{cases} 2x + y = 2 \\ 3x + 7y = 14 \end{cases}$$

$$32) \begin{cases} x = 3y + 4 \\ 2x - 6y = 8 \end{cases}$$

$$33) \begin{cases} 4x - y = 6 \\ y = 4x - 3 \end{cases}$$

$$34) \begin{cases} y = x + 7 \\ x - y = 3 \end{cases}$$

$$35) \begin{cases} -9x + 6y = 21 \\ 2y - 7 = 3x \end{cases}$$

$$36) \begin{cases} y = 3x + 5 \\ y = 3x - 1 \end{cases}$$

$$37) \begin{cases} 3x - 3y = -24 \\ x - y = 5 \end{cases}$$

$$38) \begin{cases} x + 2y = 1 \\ 5x = 5 - 10y \end{cases}$$

$$39) \begin{cases} 12x + 6y = 0 \\ y = -2x \end{cases}$$

$$40) \begin{cases} -x - 4y = -14 \\ -6x + 8y = 12 \end{cases}$$

2.2 Answers

- 1) $(1, -3)$
- 2) $(-3, 2)$
- 3) $(-2, -5)$
- 4) $(0, 3)$
- 5) $(-1, -2)$
- 6) $(-7, -8)$
- 7) $(1, 5)$
- 8) $(-4, -1)$
- 9) $(3, 3)$
- 10) $(4, 4)$
- 11) $(2, 6)$
- 12) $(-3, 3)$
- 13) $(-2, -6)$
- 14) $(0, 2)$
- 15) $(1, -5)$
- 16) $(-1, 0)$
- 17) $(-1, 8)$
- 18) $(3, 7)$
- 19) $(2, 3)$
- 20) $(8, -8)$
- 21) $(\frac{4}{5}, \frac{21}{5})$
- 22) $(\frac{5}{7}, \frac{53}{7})$
- 23) $(-3, -2)$
- 24) $(1, -3)$
- 25) $(-\frac{25}{7}, \frac{17}{7})$
- 26) $(\frac{78}{31}, \frac{7}{31})$
- 27) $(-2, 8)$
- 28) $(-4, 3)$
- 29) $(4, -3)$
- 30) $(-1, 5)$
- 31) $(0, 2)$
- 32) Infinite Solutions
- 33) No Solution

- 34) No Solution
- 35) Infinite Solutions
- 36) No Solution
- 37) No Solution
- 38) Infinite Solutions
- 39) Infinite Solutions
- 40) (2, 3)